

## The Role of Er YAG Laser in Modification of Pigmentation of Scar

Nishad K<sup>1</sup>, Ravi Kumar Chittoria<sup>2</sup>, Padmalakshmi Bharathi Mohan<sup>3</sup>,  
Imran Pathan<sup>4</sup>, Shijina K<sup>5</sup>, Neljo Thomas<sup>6</sup>

### Abstract

The scar is a sequela of any trauma, surgery, or burns. If the scar becomes unsightly it influences the psychological or physical well beingness of the patient. The pigmentation difference of the scar with the surrounding skin is one factor that can make the scar unsightly. There are many methods of scar management. Laser Therapy using Er YAG laser is one method of management of scars that are widely used in western countries. In India, the data on scar management using Er YAG laser and the effect of Er YAG laser on the pigmentation of the scar is few. This study was conducted in a tertiary care center in South India on 73 scars for 6 months

**Keywords:** Scar management; Scar Pigmentation; Laser Therapy; Er-YAG Laser.

### Introduction

Scar management is a very common problem for which patients seek consultation from any plastic surgeon. Abnormal scars can be painful, itchy and it can disable the patient as scars it can prevent the movement of the joint, neck eyelid, or lips. Scars can become unsightly due to their location, color, consistency, or size (height). There are many methods for the prevention and management of scars, Though the scars are not completely avoidable, they can certainly be made better with meticulous management of the wounds. There is no single best method to manage the scar. Some of the commonly

used methods of scar management are scar massage with emollients, compressions garments, intralesional steroids, surgical scar revision, and Laser Therapy. Laser therapy can be low level Laser Therapy as well as high level Laser therapy.

High level laser therapy is in use for many years, initial lasers in use were CO<sub>2</sub> and Pulsed dye laser. Due to the adverse effects, there is always a quest for newer and newer lasers that are equally effective and with lesser adverse effects and lesser downtime to achieve the desired clinical change. The desired clinical changes that can make a scar less unsightly are changes in size (height), consistency, color (pigmentation), vascularity.

Though the Er-YAG laser was used in western countries for many years, in India Er YAG is a relatively newer addition to the armamentarium in scar management, hence the data of the efficacy of the Er YAG in managing unsightly scars in Indian skin type is few. In this study, we have used the Er YAG for fractional ablative resurfacing of the post-trauma and burns scars and studied the effect of the Laser on each scar parameter. The efficacy of the Laser on pigmentation is studied in special interest as pigmentation is one parameter that can attract a beholder's attention to any scar.

### Materials and Methods

This study was conducted in the Department of Plastic Surgery at a tertiary care center after getting the departmental ethical committee approval. Informed written consent was taken from the patients for Er YAG therapy as well as the clinical photography. The details of the study are as follows: 73 scars were enrolled into the study randomly which included post-trauma, post-surgical well as the post burns scars. The scars were evaluated twice using the Vancouver scar scale scoring system and clinical photography twice once pre-treatment and next one month after the completion of the laser therapy. The laser therapy was given for four sessions each at a one-month interval. The laser used was Er: YAG Laser, Twain 2940, Quanta System S.p.A., Italy, in ablative as well as thermal mode, at a wavelength of

### Authors Affiliation

<sup>1</sup>Senior Resident, <sup>2</sup>Professor, <sup>3-6</sup>Resident, Department of Plastic Surgery, Jawaharlal Institute of Postgraduate Medical Education and Research, Pondicherry 605006, India.

### Corresponding Affiliation

**Ravi Kumar Chittoria**, Professor, Department of Plastic Surgery, Jawaharlal Institute of Postgraduate Medical Education and Research, Pondicherry 605006, India.

**Email:** drchittoria@yahoo.com

### How to cite this article:

Nishad K, Ravi Kumar Chittoria, Padmalakshmi Bharathi Mohan et al./The Role of Er YAG Laser in Modification of Pigmentation of Scar 2021;8(1):17-19.

2,940nm, fluence was set to 1 to 2 J/cm<sup>2</sup>, pulse width used was 300 microseconds using spot diameter of 4mm. During each session, two laser passes of 400 mJ in short pulse mode (pulse duration 0.30ms) and one pass of 800 mJ in long pulse mode (pulse duration 1 ms) were performed.


Statistical analysis was performed using IBM statistical software, SPSS Statistics version 27 (IBM Inc.). Normally distributed data were expressed as mean  $\pm$ SD. Data were expressed as median (interquartile range, IQR), when the assumption of normality was violated (Shapiro Wilk test,  $P < 0.001$ ). Paired T-Test was used and wherever needed one-way repeated measure ANOVA was done to determine whether there are any statistically significant differences between the means of three or more levels of a within-subjects factor over time. A  $P$  value  $< 0.05$  was considered statistically significant

## Result

The mean age of patients was  $35.2 \pm 7.8$  (range, 18-50 years), respectively. There were 43 males with a male-female ratio of 2.1:1. The preprocedural and postprocedural Vancouver scar scale parameters are comparisons showed that there was a significant difference in pigmentation, after laser application (paired t-test,  $P = 0.006$ )

**Table 1:** Preprocedural and Post Procedural Changes of Vancouver scar scale variables.

VSS Parameter	Pre Procedural	Post Procedural	P Value
Vascularity	1.4+0.9	1.9+0.8	0.001
Pigmentation	1.47+0.6	1.2+0.6	0.006
Pliability	2.1+0.8	1.5+0.6	0.001
Height	1.6+0.7	1+0.6	0.001
Total Score	6.6+1.6	5.6+1.4	0.001



**Fig. 2:** Graph Showing the changes of VSS variables.

## Discussion

The scar is defined as fibrous tissue that replaces the wound after injury or disease.<sup>2</sup> During the process of healing the wound develops a bridge of collagen fibers with a thin epithelium, forming an immature scar.<sup>3</sup> An immature scar is red, raised, rigid, and hypopigmented, During the process of maturation the scar becomes more pliable, flatter, less vascular color is normalized. Any aberration of these makes the scar abnormal or unsightly. The difference between the normal scar, immature scar hypertrophic scar lies in the difference in the extracellular matrix composition, Type-III is the predominant type of collagen during the proliferation phase of normal wound healing and which eventually gets replaced with type-I collagen during the remodeling phase. A normal scar when mature consists of 80% type-I collagen with 10-15% type-III and a minimal amount of type-V collagen. This composition is altered in an abnormal scar with an increased ratio of type-III to type-I collagen. The abnormal scar consists of around 33% type-III, 10% type-V, and around 60% type-I collagen. Apart from the composition of the collagen, the arrangement of fibrils and interfibrillar space also is different in an abnormal scar compared to the normal mature scar. The cellular function of fibroblasts and keratinocytes is also altered in an abnormal scar making them profibrotic. The expression of cytokines is also altered in an abnormal scar.

The balance between matrix metalloproteinase (MMPs) and tissue inhibitors of metalloproteinase (TIMPs) is altered and is moved towards the profibrotic side. Transforming growth factor- $\beta$  (TGF- $\beta$ ), connective tissue growth factor (CTGF), platelet-derived growth factor (PDGF), and insulin-like growth factor 1 (ILGF- $\alpha$ ) are up-regulated, meanwhile interferon  $\alpha$  (IFN- $\alpha$ ) and interferon- $\gamma$  (IFN- $\gamma$ ) are down regulated.<sup>4,5</sup>

The first LASER machine was devised in 1960 by Maimon, which was a Ruby laser. Dr. Leon Goldman a dermatologist is considered to be the father of laser medicine. The first laser that was specifically designed for use in a medical condition was Pulsed Dye Laser (PDL), which was used for port-wine stains. Since then laser technology has evolved a lot with newer concepts of pulsed therapy, fractionated laser therapy, Q-switched mode, etc being added to the list. The principle of any laser is photo thermolysis, which was proposed first by Anderson. Each laser has a specific target on which it acts, known as chromophore. The laser selectively acts on its chromophore and produces thermal ablation

of the target tissue. Fluence, pulse width, spot size, and stacking are variables that are to be adjusted according to the individual requirements. The mechanism by which a laser affects scar remodeling is not fully known, but ablative fractional resurfacing may lead to the production of various cytokines and growth factors by stimulating a variety of not fully known cellular responses. Fractional photothermolysis produces controlled and limited dermal heating which triggers a cascade of events in which normalization of the collagenesis-collagenolysis cycle occurs.

### Conclusion

The study shows that Er YAG Laser therapy is an effective method in the management of post-trauma scar. The pigmentation of the scar showed significant improvement after the application of the Er YAG Laser. No adverse effects were noted during the study. The limitation of the study is that there is no separate comparison for hypopigmentation and hyperpigmentation after the Laser therapy. We suggest a study for hypopigmentation and

hyperpigmentation separately. Large volume and multi-center study may give a better picture of the effect of Er YAG laser.

### Competing interest

Declarations: None

Author's contributions: All authors made contributions to the article

Financial support and sponsorship: None

### References

1. Stedman TL, ed. Stedman's Medical Dictionary. 23rd ed. Baltimore, MD: Williams and Wilkins; 1976.
2. Serghiou MA, Ott S, Cowan A, Offenbergl JK, Suman OE. Burn Rehabilitation Along the Continuum of Care. In: Herndon DN, editor. Total Burn Care. 5th ed. Edinburgh: Elsevier; 2018: 490-495.
3. Kwan P, Desmouliere A, Tredget EE. Molecular and Cellular basis of Hypertrophic Scarring. In: Herndon DN, editor. Total Burn Care. 5th ed. Edinburgh: Elsevier; 2018: 455-465.
4. Hawkins HK, Jay J, Finnerty CC. Pathophysiology of the Burn Scar. In: Herndon DN, editor. Total Burn Care. 5th ed. Edinburgh: Elsevier; 2018. p. 466-475.



# REDKART.NET

(A product of Red Flower Publication (P) Limited)  
(Publications available for purchase: Journals, Books, Articles and Single issues)  
(Date range: 1967 to till date)

The Red Kart is an e-commerce and is a product of Red Flower Publication (P) Limited. It covers a broad range of journals, Books, Articles, Single issues (Print & Online-PDF) in English and Hindi languages. All these publications are in stock for immediate shipping and online access in case of online.

**Benefits of shopping online are better than conventional way of buying.**

1. Convenience.
2. Better prices.
3. More variety.
4. Fewer expenses.
5. No crowds.
6. Less compulsive shopping.
7. Buying old or unused items at lower prices.
8. Discreet purchases are easier.

URL: [www.redkart.net](http://www.redkart.net)